

Running charm quark mass versus M_D and $D_{(s)} \rightarrow \mu\nu$

Stephan Narison

Laboratoire de Physique Mathématique et Théorique
UM2, Place Eugène Bataillon
34095 Montpellier Cedex 05, France
E-mail: narison@lpm.univ-montp2.fr

Abstract

I study (for the first time) the dependence of M_D and the leptonic decay constant f_D on the variation of the running charm quark mass $\bar{m}_c(\nu)$. I conclude that the present data on f_{D_s} from $D_s \rightarrow \mu\nu$ decay give a weaker constraint than M_D , where the latter leads to the result: $\bar{m}_c(M_c) = (1.08 \pm 0.11)$ GeV, to two-loop accuracy in the $\overline{MS}$ scheme, in agreement with the value $\bar{m}_c(M_c) = (1.23^{+0.04}_{-0.05})$ GeV extracted directly, within the same approximation, from $M_{J/\psi}$. The agreement of the corresponding perturbative pole mass with the one extracted directly from the data can indicate that the non-perturbative effects to the pole mass are negligible. Inversely, injecting the average value of $\bar{m}_c(M_c)$ from M_D and $M_{J/\psi}$ into the $\bar{m}_c$ behaviour of f_D , I obtain $f_D \simeq (1.52 \pm 0.16)f_\pi$, which combined with the sum rule prediction for f_{D_s}/f_D , gives $f_{D_s} \simeq (1.75 \pm 0.18)f_\pi$ in good agreement with the data. The extension of the analysis to the case of f_B and f_{D^*} is discussed.

1 Introduction

One of the most important parameters of the standard model is the quark masses. However, contrary to the leptons, where the physical mass can be identified with the pole of the propagator, the quark masses are difficult to define because of confinement. Some attempts have been made in order to define the quark pole mass within perturbation theory, where it has been shown to be IR-finite [1] and independent from the choice of the regularization and renormalization schemes used [2]. More recently, it has been noticed, in the limit of a large number of flavours, that the resummation of perturbative series can induce a non-perturbative term, which can affect the truncated perturbative result, and can, then, limit the accuracy of the pole mass determination [3]. However, a proper use of such a result needs a resummation, at the same level of accuracy (which is not often the case), of the Green function of a given process involving the pole mass, where some eventual cancellation between the resummed terms of the two series may occur. One may bypass the previous problems, by working, at a given order of perturbative QCD, with the running quark masses, which are treated like coupling constants of the QCD Lagrangian (see e.g. [4]), and where some non-perturbative-like effect is expected to be absent. A lot of efforts has been furnished in the literature [5] for extracting directly from the data the running masses of the light and heavy quarks using the SVZ QCD spectral sum rules (QSSR) [6, 7]. In this note, I shall consider a direct extraction of the running charm quark mass using the observed value of $M_D = 1.865$ MeV and the new data on f_{D_s} from $D_s \rightarrow \mu\nu$ decay, where its average value is [8]:

$$f_{D_s} \simeq (1.92 \pm 0.23)f_\pi. \quad (1)$$

For convenience, we shall also use the fact that $r \equiv f_{D_s}/f_D$ is under good control from different non-perturbative approaches. For definiteness and for an internal self-consistency of the analysis, we shall use the QSSR value [9]:

$$\frac{f_{D_s}}{f_D} = 1.15 \pm 0.04 \implies f_D \simeq (1.67 \pm 0.24)f_\pi. \quad (2)$$

We shall extend the analysis to the case of f_B and f_{D^*} .

2 The QCD spectral sum rules

We shall work with the pseudoscalar two-point correlator:

$$\psi_5(q^2) \equiv i \int d^4x e^{iqx} \langle 0 | T J_q(x) J_q^\dagger(0) | 0 \rangle, \quad (3)$$

built from the heavy-light quark current: $J_d(x) = (m_c + m_d)\bar{c}(i\gamma_5)d$, and which has the quantum numbers of the D meson. The corresponding Laplace transform sum rules are:

$$\mathcal{L}(\tau) = \int_{t_\leq}^{\infty} dt e^{-t\tau} \frac{1}{\pi} \text{Im}\psi_5(t), \quad \text{and} \quad \mathcal{R}(\tau) \equiv -\frac{d}{d\tau} \log \mathcal{L}(\tau), \quad (4)$$

where $t_\leq$ is the hadronic threshold. The latter sum rule, or its slight modification, is also useful, as it is equal to the resonance mass squared, in the simple duality ansatz parametrization of the spectral function:

$$\frac{1}{\pi} \text{Im}\psi_5(t) \simeq 2f_D^2 M_D^4 \delta(t - M_D^2) + \text{“QCD continuum”} \Theta(t - t_c), \quad (5)$$

where the “QCD continuum comes from the discontinuity of the QCD diagrams, which is expected to give a good smearing of the different radial excitations¹. The decay constant f_D is analogous to $f_\pi = 93.3$ MeV; t_c is the QCD continuum threshold, which is, like the sum rule variable τ , an a priori arbitrary parameter. In this paper, we shall impose the t_c and τ stability criteria for extracting our optimal results². The QCD expression of the correlator is known to two-loop accuracy (see e.g. [7] and the explicit expressions given in [10]), in terms of the perturbative pole mass M_c , and including the non-perturbative condensates of dimensions less than or equal to six³. The sum rule reads:

$$\mathcal{L}(\tau) = M_c^2 \left\{ \int_{4M_c^2}^{\infty} dt e^{-t\tau} \frac{3}{8\pi^2} t(1-x)^2 \left[1 + \frac{4}{3} \left(\frac{\alpha_s}{\pi} \right) f(x) \right] + [C_4 \langle O_4 \rangle + C_6 \langle O_6 \rangle \tau] e^{-M_c^2 \tau} \right\}, \quad (6)$$

¹At the optimization scale, its effect is negligible, such that a more involved parametrization is not necessary.

²The corresponding t_c value very roughly indicates the position of the next radial excitations.

³We shall use the corrected coefficient of the quark-gluon mixed condensate given in [11]. This change affects only slightly the result. We shall also skip the negligible contribution from the dimension six four-quark and three-gluon condensates. Notice that there is some discrepancy on the value of the four-quark coefficient in the literature.

with:

$$\begin{aligned}
x &\equiv M_c^2/t, \\
f(x) &= \frac{9}{4} + 2\text{Li}_2(x) + \log x \log(1-x) - \frac{3}{2} \log(1/x-1) \\
&\quad - \log(1-x) + x \log(1/x-1) - (x/(1-x)) \log x, \\
C_4\langle O_4 \rangle &= -M_c \langle \bar{d}d \rangle - \langle \alpha_s G^2 \rangle / 12\pi \\
C_6\langle O_6 \rangle &= \frac{M_c^3 \tau}{2} g \langle \bar{d} \sigma_{\mu\nu} \frac{\lambda_a}{2} G_a^{\mu\nu} d \rangle
\end{aligned} \tag{7}$$

It can be expressed in terms of the running mass $\bar{m}_c(\nu)$ ⁴, through the one-loop relation [2]:

$$M_c(\nu) = \bar{m}_c(\nu) \left\{ 1 + \left(\frac{\bar{\alpha}_s}{\pi} \right) \left(\frac{4}{3} + 2 \log \frac{\nu}{M_c} \right) + \dots \right\}. \tag{8}$$

Throughout this paper we shall use the values of the parameters [7, 12] given in Table 1. We have used for the mixed condensate the parametrization:

$$g \langle \bar{d} \sigma_{\mu\nu} \frac{\lambda_a}{2} G_a^{\mu\nu} d \rangle = M_0^2 \langle \bar{d}d \rangle. \tag{9}$$

We shall also use, for four active flavours [13, 5]:

$$\Lambda = (325 \pm 100) \text{ MeV}. \tag{10}$$

One can inspect that the dominant non-perturbative contribution is due to the dimension-four $m_c \langle \bar{d}d \rangle$ light quark condensate, while the other non-perturbative effects remain a small correction at the optimization scale, which corresponds to $\tau \simeq 0.6 \sim 1 \text{ GeV}^{-2}$ and $t_c \simeq 6 \sim 8 \text{ GeV}^2$.

3 Discussions and results




Figure 1: Behaviour of f_D versus $\bar{m}_c(\nu)$. The horizontal band is the experimental domain of f_D . The theoretical band is limited by the two curves: (a) $\Lambda=0.425 \text{ GeV}$, $\nu=1 \text{ GeV} \approx \tau^{-1/2}$, $t_c \geq 7 \text{ GeV}^2$, $\langle \bar{d}d \rangle^{1/3}(1 \text{ GeV}) = -238 \text{ MeV}$, $M_0^2 = 0.7 \text{ GeV}^2$, $\langle \alpha_s G^2 \rangle = 0.06 \text{ GeV}^2$ and (b) $\Lambda=0.225 \text{ GeV}$, $\nu \approx M_c = 1.42 \text{ GeV}$, $t_c = 6 \text{ GeV}^2$, $\langle \bar{d}d \rangle^{1/3}(1 \text{ GeV}) = -220 \text{ MeV}$, $M_0^2 = 0.6 \text{ GeV}^2$, $\langle \alpha_s G^2 \rangle = 0.08 \text{ GeV}^2$.

Given the experimental value on $M_D = 1.865 \text{ GeV}$, we present our results on f_D from the first sum rule for different values of the charm quark running mass evaluated at $p^2 = \nu^2$ in Fig. 1. The second

⁴It is clear that, for the non-perturbative terms which are known to leading order of perturbation theory, one can use either the running or the pole mass. However, we shall see that the non-perturbative effects are not important in the analysis, such that this distinction does not affect the result.

sum rule gives the prediction on M_D for each value of the charm mass (Fig. 2). The theoretical band is limited by the two extremal values of the QCD parameters used in Table 1. Notice that the effect of the errors of the different input parameters is much smaller in the ratio of sum rule $\mathcal{R}(\tau)$. The previous analysis leads to the prediction from a two-loop calculation in the $\overline{MS}$ scheme:

$$\bar{m}_c(M_c) = (1.08 \pm 0.11) \text{ GeV}, \quad (11)$$

where one should notice that, within the present accuracy of the data on f_{D_s} , the result comes mainly from the one used to reproduce the experimental value of M_D .




Figure 2: Behaviour of M_D versus $\bar{m}_c(\nu)$. The horizontal band is the experimental value of M_D . The theoretical band is limited by the two curves: (a) the same as (a) of Fig. 1 but $\nu=1$ GeV and (b) the same as (b) of Fig. 1 but $\nu=1.42$ GeV.

We have varied the subtraction scale ν in the range from $\tau^{-1/2} \approx 1$ GeV to $M_c \approx 1.42$ GeV. One should also notice, in Table 1, that the effect of t_c on the result is relatively small from the value $t_c \simeq 6 \text{ GeV}^2$, where one starts to have a τ stability until $t_c \geq 7 \text{ GeV}^2$, where one has t_c stability. The main sources of errors are due to Λ and ν . Within the errors, the present result is in good agreement (though less accurate) with the value [14]:

$$\bar{m}_c(M_c) = (1.23^{+0.04}_{-0.05}) \text{ GeV} \quad (12)$$

obtained, within the same two-loop approximation, from $M_{J/\psi}$. Inversely, we can use the combined value:

$$\bar{m}_c(M_c) = (1.20 \pm 0.05) \text{ GeV} \quad (13)$$

from M_D and from $M_{J/\psi}$ systems on the curve f_D as function of $\bar{m}_c$ given in Fig. 1. Then, one can deduce:

$$f_D \simeq (1.52 \pm 0.16) f_\pi, \quad (14)$$

where, as can be seen in Table 1, the errors in this determination come mainly from the perturbative parameters: Λ , m_c and the subtraction scale ν , and, to a lesser extent, from t_c and the non-perturbative terms. We can compare this result with the previous value:

$$f_D \simeq (1.35 \pm 0.07) f_\pi, \quad (15)$$

obtained by using the perturbative pole mass of the charm quark propagator [9, 10]. The good

Table 1: Different sources of errors in the estimate of f_D

Sources	$ \Delta(f_D/f_\pi) $
$\Lambda = (325 \pm 100) \text{ MeV}$	0.12
$\nu = (1.20 \pm 0.22) \text{ GeV}$	0.08
$\bar{m}_c(M_c) = (1.20 \pm 0.05) \text{ GeV}$	0.05
$t_c = (6.5 \pm 0.5) \text{ GeV}^2$	0.04
$\langle \bar{d}d \rangle^{1/3}(1 \text{ GeV}) = -(229 \pm 9) \text{ MeV}$	0.02
$M_0^2 = (0.8 \pm 0.1) \text{ GeV}^2$	0.02
$\langle \alpha_s G^2 \rangle = (0.07 \pm 0.01) \text{ GeV}^2$	0.01
Total	0.16

agreement between the two results in Eqs (14) and (15) within the errors may be an indirect indication that the pole mass defined at a given order of perturbation theory, can provide a good description of the physical process in this channel, and that the eventual non-perturbative power corrections induced by the resummation of the perturbative series remain small corrections. This observation can also be supported by the agreement of the value of the perturbative pole mass obtained here and of the one from the J/ψ systems, where both values have been obtained at two-loop accuracy. A further support of this argument can also be provided by the agreement of the pole mass deduced from Eq. (8) using the value of the running mass, with the one extracted directly in [14]. Finally, using the previous value of the ratio f_{D_s}/f_D given in Eq. (2), one obtains:

$$f_{D_s} \simeq (1.75 \pm 0.18)f_\pi, \quad (16)$$

which is in good agreement with the data quoted in Eq. (1). A natural improvement of the analysis done in this paper is a much more precise measurement of f_{D_s} or/and an evaluation of the QCD two-point correlator to three-loop accuracy. The two projects are expected to be feasible in the near future.

4 Extension of the analysis to f_B and f_{D^*}

The extension of the previous analysis to the case of the B meson is not very conclusive due to the strong dependence of the result on the value of the subtraction scale ν , which can vary, for the B meson case, in a larger range. Instead, using our previous observation on the agreement of the results on f_D from the running and perturbative pole masses, we also expect that the result obtained from the pole mass of the b quark [9, 10] should also be reliable, which is:

$$f_B \simeq (1.49 \pm 0.08)f_\pi. \quad (17)$$

We have also extended the previous analysis to the case of the vector current having the quantum number of the D^* . In this case, and working with the correlator having the same dimension as $\psi_5(q^2)$ ⁵, we do not obtain a τ stability as the coefficients of the chiral condensates are opposite of the pseudoscalar ones. Then, we conclude that, from this quantity, we cannot have a good determination of f_{D^*} .

Acknowledgements

I thank, for its hospitality, the CERN Theory Division, where this work has been finalized.

⁵Notice that, in this channel, the correlator which has less power of q^2 than $\psi_5(q^2)$, and which might present a τ stability, can have singularities and then should be used with a great care.

References

- [1] R. Tarrach, *Nucl. Phys.* **B183** (1981) 384;
- [2] S. Narison, *Phys. Lett.* **B197** (1987) 405; **B216** (1989) 191.
- [3] M. Beneke and V.M. Braun, *Nucl. Phys.* **B426** (1994) 301; P. Ball, M. Beneke and V.M. Braun, *Nucl. Phys.* **B452** (1995) 563; I. Bigi et al., *Phys. Rev.* **D50** (1994) 2234; M. Beneke (private communication); G. Grunberg (private communication); For a review, see e.g.: R. Akhoury and V. I. Zakharov, *Nucl. Phys. (Proc. Suppl.)* **B, A54** (1997); *QCD97*, Montpellier (1997) and references therein.
- [4] E.G. Floratos, S. Narison and E. de Rafael, *Nucl. Phys.* **B155** (1979) 115.
- [5] PDG96, R.M. Barnett et al., *Phys. Rev.* **D54** (1996) 1.
- [6] M.A. Shifman, A.I. Vainshtein and V.I. Zakharov, *Nucl. Phys.* **B147** (1979) 385, 448.
- [7] For a review and references to original works, see e.g., S. Narison, *QCD spectral sum rules, Lecture Notes in Physics*, **Vol 26** (1989) ed. World Scientific; *Acta Phys. Pol.* **26** (1995) 687; *Recent Progress in QCD spectral sum rules* (to appear).
- [8] J.D. Richman, *ICHEP96*, Warsaw (1996).
- [9] S. Narison, *Phys. Lett.* **B322** (1994) 247.
- [10] S. Narison, *Phys. Lett.* **B198** (1987) 104; **B210** (1988) 238.
- [11] S. Generalis, *J. Phys., Nucl. Part. Phys.* **G16** (1990) 367.
- [12] S. Narison, *Phys. Lett.* **B361** (1995) 121; **B387** (1996) 162.
- [13] S. Bethke, *Nucl. Phys. (Proc. Suppl.)* **B, C39** (1995); *ibid*, **B, A54** (1997); I. Hinchliffe, *Meeting of the American Physical Society*, Albuquerque, 1994; M. Schmelling, *ICHEP96*, Varsaw, 1996.
- [14] S. Narison, *Phys. Lett.* **B341** (1994) 73.